

Source: Centers for Disease Control and Prevention (CDC)



Developed by IPRO ESRD Network of the Ohio River Valley while under contract with Centers for Medicare & Medicaid Services. Contract HHSM-500-2016-00009C



End-Stage Renal Disease Network of the Ohio River Valley





End-Stage Renal Disease Network of the Ohio River Valley

Protect yourself.

Get the Vaccines You Need!

Vaccination is a safe, effective way to protect yourself from serious illness.

For more information or to file a grievance, please contact us:

End-Stage Renal Disease Network of the Ohio River Valley 3201 Enterprise Parkway, Suite 210 Beachwood, OH 44122

Toll-Free Patient Phone: (844) 819-3010

Main Phone: (216) 593-0001

Fax: (216) 593-0101

E-mail: esrdnetwork9@ipro.us

website: esrd.ipro.org

Vaccines recommended for dialysis patients:

- Annual Flu Vaccine
- Pneumonia Vaccine
- Hepatitis Vaccine

IPRO End-Stage Renal Disease Network of the Ohio River Valley (844) 819-3010 esrd.ipro.org

Annual Influenza (Flu) Vaccine



- Influenza, also called the flu, is a contagious and serious respiratory disease.
- As a dialysis patient, if you get the flu is soon are more likely than others to develop serious problems:
- Each year there are different types of flu-vaccines available; some are better suited for kidney patients.

 Ask your healthcare team about which flu vaccine is best for you.
- Receiving an annual flu vaccine will help protect you from getting the flu
- According to the Centers for Disease Control and Prevention (CDC), influenza season usually is at its worst in February and can last until late May. The best time to receive a vaccine is October or November.

Pneumonia Vaccine

Prieumonia, an infection of the lungs needlessly affects millions of people worldwide each year.

Phaumonia is caused by bacteria and

Prieumonia infections can often be prevented and can usually be treated

 The pneumonia racine protects you body from many types of Harmful bacteria

You should receive a proumontal vaccine every five years.

ou can receive this vaccine any time



Hepatitis Va**c**ine

inflamed, and inflamed, and inflamed, and inflamed, and inflamed incomed inflamed infections.



even life threatening.

lepaini. Bit spread through contains to look or body fluids from smeare who has the virus.

Italysis patients are at greater risk or exposure to this virus because of epeared access to the bloodstream luring steetment.

caused by hepatitis B.

 The hepatitis B vaccine is usually given in a series of three to four injections or doses over a six-month period.

For more information about these vaccines, speak with your healthcare team and visit these CDC website pages:

www.cdc.gov/flu • www.cdc.gov/pneumococcal/vaccination.html • www.cdc.gov/hepatitis/abc